

REFERENCES

- Alexander, E.C., Davis, M.A., and Alexander, S.C., 1989, Hydrologic study of Jewel Cave/Wind Cave, Final Report: University of Minnesota, unpublished report, contract CX-1200-S-A047, 149 p.
- Alexander, E.C., Davis, M.A., Alexander, S.C., and Lively, R.S., 1988, Thermal springs of the southern Black Hills, *in* Schilberg and Springhetti, eds., Caves and associated features of the Black Hills: Guidebook 45th Annual Convention National Speleological Society, p. 14-26.
- Anderson, M.T., Driscoll, D.G., and Williamson, J.E., 1999, Ground-water and surface-water interactions along Rapid Creek near Rapid City, South Dakota: U.S. Geological Survey Water-Resources Investigations Report 98-4214, 99 p.
- Back, William, Hanshaw, B.B., Plummer, L.N., Rahn, P.H., Rightmire, C.T., and Rubin, Meyer, 1983, Process and rate of dedolomitization; mass transfer and ¹⁴C dating in a regional carbonate aquifer: Geological Society of America Bulletin, v. 94, p. 1415-1429.
- Bowles, C.G., and Braddock, W.A., 1963, Solution breccias of the Minnelusa Formation in the Black Hills, South Dakota and Wyoming: U.S. Geological Survey Professional Paper 475-C, p. C91-C95.
- Braddock, W.A., 1963, Geology of the Jewel Cave SW quadrangle, Custer County, South Dakota: U.S. Geological Survey Bulletin 1063-G, p. 217-268.
- Braddock, W.A., and Bowles, C.G., 1963, Calcitization of dolomite by calcium sulfate solutions in the Minnelusa Formation, Black Hills, South Dakota and Wyoming: U.S. Geological Survey Professional Paper 475-C, p. 96-99.
- Brobst, D.A., and Epstein, J.B., 1963, Geology of the Fanny Peak quadrangle, Wyoming-South Dakota: U.S. Geological Survey Bulletin 1063-I, p. 323-377.
- Browne, G.S., 1992, Water quality characteristics and geochemical differentiation of the Minnelusa and Madison aquifers in the vicinity of Rapid City, South Dakota: Rapid City, South Dakota School of Mines and Technology, unpublished M.S. thesis, 84 p.
- Busby, J.F., Kimball, B.A., Downey, J.S., and Peter, K.D., 1995, Geochemistry of water in aquifers and confining units of the Northern Great Plains in parts of Montana, North Dakota, South Dakota, and Wyoming: U.S. Geological Survey Professional Paper 1402-F, 146 p.
- Busby, J.F., Lee, R., and Hanshaw, B.B., 1983, Major geochemical processes related to the hydrology of the Madison aquifer system and associated rocks in parts of Montana, South Dakota, and Wyoming: U.S. Geological Survey Water-Resources Investigation Report 83-4093, 180 p.
- Busby, J.F., Plummer, L.N., Lee, R.W., and Hanshaw, B.B., 1991, Geochemical evolution of ground water in the Madison aquifer system in parts of Montana, Wyoming, and South Dakota: U.S. Geological Survey Professional Paper 1273-F, 89 p.
- Carter, J.M., Driscoll, D.G., and Hamade, G.R., 2001, Estimated recharge to the Madison and Minnelusa aquifers in the Black Hills area, South Dakota: U.S. Geological Survey Water-Resources Investigations Report 00-4278, 66 p.
- Carter, J.M., Driscoll, D.G., Hamade, G.R., and Jarrell, G.J., 2001, Hydrologic budgets for the Madison and Minnelusa aquifers, Black Hills of South Dakota and Wyoming, water years 1987-96: U.S. Geological Survey Water-Resources Investigations Report 01-4119, 53 p.
- Carter, J.M., and Redden, J.A., 1999a, Altitude of the top of the Inyan Kara Group, Black Hills area, South Dakota: U.S. Geological Survey Hydrologic Investigation Atlas HA-744-A, 2 sheets, scale 1:100,000.
- 1999b, Altitude of the top of the Minnekahta Limestone, Black Hills area, South Dakota: U.S. Geological Survey Hydrologic Investigation Atlas HA-744-B, 2 sheets, scale 1:100,000.
- 1999c, Altitude of the top of the Minnelusa Formation, Black Hills area, South Dakota: U.S. Geological Survey Hydrologic Investigation Atlas HA-744-C, 2 sheets, scale 1:100,000.
- 1999d, Altitude of the top of the Madison Limestone, Black Hills area, South Dakota: U.S. Geological Survey Hydrologic Investigation Atlas HA-744-D, 2 sheets, scale 1:100,000.
- 1999e, Altitude of the top of the Deadwood Formation, Black Hills area, South Dakota: U.S. Geological Survey Hydrologic Investigation Atlas HA-744-E, 2 sheets, scale 1:100,000.
- Clark, I.D., and Fritz, P., 1997, Environmental isotopes in hydrogeology: Boca Raton, Fla., CRC Press/Lewis Publishers, 328 p.
- Craig, H., 1961, Isotopic variations in meteoric waters: Science, v. 133, p. 1702-1703.
- DeWitt, Ed, Redden, J.A., Buscher, David, and Wilson, A.B., 1989, Geologic map of the Black Hills area, South Dakota and Wyoming: U.S. Geological Survey Map I-1910, 1 sheet, scale 1:250,000.
- DeWitt, Ed, Redden, J.A., Wilson, A.B., and Buscher, David, 1986, Mineral resource potential and geology of the Black Hills National Forest, South Dakota and Wyoming: U.S. Geological Survey Bulletin 1580, 135 p.
- Downey, J.S., 1984, Geohydrology of the Madison and associated aquifers in parts of Montana, North Dakota, South Dakota, and Wyoming: U.S. Geological Survey Professional Paper 1273-G, 47 p.

- Downey, J.S., and Dinwiddie, G.A., 1988, The regional aquifer system underlying the Northern Great Plains in parts of Montana, North Dakota, South Dakota, and Wyoming—Summary: U.S. Geological Survey Professional Paper 1402-A, 64 p.
- Driscoll, D.G., 1992, Plan of study for the Black Hills Hydrology Study, South Dakota: U.S. Geological Survey Open-File Report 92-84, 10 p.
- Driscoll, D.G., Bradford, W.L., and Moran, M.J., 2000, Selected hydrologic data, through water year 1998, Black Hills Hydrology Study, South Dakota: U.S. Geological Survey Open-File Report 00-70, 284 p.
- Driscoll, D.G., and Carter, J.M., in press, Hydrologic conditions and budgets for the Black Hills of South Dakota, through water year 1998: U.S. Geological Survey Water-Resources Investigations Report 01-4226.
- Driscoll, D.G., Hamade, G.R., and Kenner, S.J., 2000, Summary of precipitation data for the Black Hills area of South Dakota, water years 1931-98: U.S. Geological Survey Open-File Report 00-329, 151 p.
- Epstein, J.B., 2000, Gypsum karst and hydrologic evolution in the northern Black Hills, South Dakota, *in* Strobel, M.L., Davis, A.D., Sawyer, J.F., Webb, C.J., Naus, C.A., and Rahn, P.H., eds., Proceedings of the 1999 Conference on the Hydrology of the Black Hills: Rapid City, South Dakota School of Mines and Technology Bulletin 20, p. 73-79.
- Fahrenbach, M.D., 1995, Sedimentation, paleontology, and paleoecology of the Late Devonian-Early Mississippian Englewood Formation, Black Hills, South Dakota: Rapid City, South Dakota School of Mines and Technology, unpublished Ph.D. dissertation, 615 p.
- Feldman, R.M., and Heimlich, R.A., 1980, The Black Hills: Kent, Ohio, Kent State University, Kendall/Hunt Publishing Company, K/H Geology Field Guide Series, 190 p.
- Ford, D.C., Lundberg, J., Palmer, A.N., Palmer, M.V., Dreybrodt, W., and Schwarcz, H.P., 1993, Uranium-series dating of the draining of an aquifer—the example of Wind Cave, Black Hills, South Dakota: Geological Society of America Bulletin, v. 105, p. 241-250.
- Freeze, R.A., and Cherry, J.A., 1979, Groundwater: Englewood Cliffs, N.J., Prentice-Hall, 604 p.
- Gonfiantini, R., Frohlich, K., Araguas-Araguas, L., and Rozanski, K., 1998, Isotopes in groundwater hydrology, *in* Kendall, C., and McDonnell, J., eds., Isotope tracers in catchment hydrology: New York, Elsevier Publishing Company, p. 203-246.
- Gott, G.B., Wolcott, D.E., and Bowles, C.G., 1974, Stratigraphy of the Inyan Kara Group and localization of uranium deposits, southern Black Hills, South Dakota and Wyoming: U.S. Geological Survey Professional Paper 763, 57 p.
- Greene, E.A., 1993, Hydraulic properties of the Madison aquifer system in the western Rapid City area, South Dakota: U.S. Geological Survey Water-Resources Investigations Report 93-4008, 56 p.
- 1997, Tracing recharge from sinking streams over spatial dimensions of kilometers in a karst aquifer: Ground Water, v. 35, no. 5, p. 898-904.
- 1999, Characterizing recharge to wells in carbonate aquifers using environmental and artificially recharged tracers, *in* Morganwalp, D.W., and Buxton, H.T., eds., U.S. Geological Survey Toxic Substances Hydrology Program—Proceedings of the Technical Meeting, Charleston, South Carolina, March 8-12, 1999—Volume 3 of 3—Subsurface Contamination from Point Sources: U.S. Geological Survey Water-Resources Investigations Report 99-4018C, p. 803-808.
- Greene, E.A., Shapiro, A.M., and Carter, J.M., 1999, Hydrogeologic characterization of the Minnelusa and Madison aquifers near Spearfish, South Dakota: U.S. Geological Survey Water-Resources Investigations Report 98-4156, 64 p.
- Gries, J.P., 1975, Mineral and water resources of South Dakota: U.S. Congress, 94th, 1st Session, Interior and Insular Affairs Committee Print, 313 p.
- 1996, Roadside geology of South Dakota: Missoula, Mont., Mountain Press Publishing Company, 358 p.
- Hayes, T.S., 1999, Episodic sediment-discharge events in Cascade Springs, southern Black Hills, South Dakota: U.S. Geological Survey Water-Resources Investigations Report 99-4168, 34 p.
- Hem, J.D., 1992, Study and interpretation of the chemical characteristics of natural water (3d ed.): U.S. Geological Survey Water-Supply Paper 2254, 263 p.
- Hortness, J.E., and Driscoll, D.G., 1998, Streamflow losses in the Black Hills of western South Dakota: U.S. Geological Survey Water-Resources Investigations Report 98-4116, 99 p.
- Huntoon, P.W., 1985, Rejection of recharge water from Madison aquifer along eastern perimeter of Bighorn Artesian Basin, Wyoming: Ground Water, v. 23, no. 3, p. 345-353.
- International Atomic Energy Agency, 1981, Statistical treatment of environmental isotope data in precipitation: Technical Report Series No. 206, 256 p.
- 1999, Global Network for Isotopes in Precipitation—The GNIP Database, Release 3, October 1999: accessed December 9, 1999, at URL <http://www.iaea.org/programs/ri/gnip/gnipmain.html>
- Johnson, B.N., 1933, A climatological review of the Black Hills: The Black Hills Engineer, Rapid City, South Dakota School of Mines and Technology, 71 p.

- Katz, B.G., Hornsby, H.D., Bohlke, J.F., and Mokray, M.F., 1999, Sources and chronology of nitrate contamination in spring waters, Suwannee River Basin, Florida: U.S. Geological Survey Water-Resources Investigations Report 99-4252, 54 p.
- Klemp, J.A., 1995, Source aquifers for large springs in northwestern Lawrence County, South Dakota: Rapid City, South Dakota School of Mines and Technology, unpublished M.S. thesis, 175 p.
- Krauskopf, K.B., and Bird, D.K., 1995, Introduction to geochemistry (3d ed.): New York, McGraw-Hill, 647 p.
- Kyllonen, D.P., and Peter, K.D., 1987, Geohydrology and water quality of the Inyan Kara, Minnelusa, and Madison aquifers of the northern Black Hills, South Dakota and Wyoming, and Bear Lodge Mountains, Wyoming: U.S. Geological Survey Water-Resources Investigations Report 86-4158, 61 p.
- Langmuir, Donald, 1997, Aqueous environmental geochemistry: Upper Saddle River, N.J., Prentice-Hall, 600 p.
- Long, A.J., 2000, Modeling techniques for karst aquifers—anisotropy, dual porosity, and linear systems analysis: Rapid City, South Dakota School of Mines and Technology, unpublished Ph.D. dissertation, 59 p.
- Long, A.J., and Putnam, L.D., 2000, Conceptual model of the Minnelusa and Madison aquifers near Rapid City, South Dakota, *in* Strobel, M.L., Davis, A.D., Sawyer, J.F., Webb, C.J., Naus, C.A., and Rahn, P.H., eds., Proceedings of the 1999 Conference on the Hydrology of the Black Hills: Rapid City, South Dakota School of Mines and Technology Bulletin no. 20, p. 98-104.
- Long, A.J., Strobel, M.L., and Hamade, G.H., 1999, Localized leakage between the Madison and Minnelusa aquifers in the eastern part of the Black Hills, South Dakota [abs.]: Geological Society of America Abstracts with Programs, October 25-28, 1999, p. 412.
- Maloszewski, P., and Zuber, A., 1982, Determining the turnover time of groundwater systems with the aid of environmental tracers, 1. Models and their applicability: *Journal of Hydrology*, v. 57, p. 207-231.
- Manga, M., 1999, On the timescales characterizing groundwater discharge at springs: *Journal of Hydrology*, v. 219, p. 56-69.
- Michel, R.L., 1989, Tritium deposition in the continental United States, 1953-83: U.S. Geological Survey Water-Resources Investigations Report 89-4072, 46 p.
- Miller, L.D., and Driscoll, D.G., 1998, Streamflow characteristics for the Black Hills of South Dakota, through water year 1993: U.S. Geological Survey Water-Resources Investigations Report 97-4288, 322 p.
- National Atmospheric Deposition Program/National Trends Network, 1999, National Atmospheric Deposition Program/National Trends Network monitoring location WY99: accessed February 12, 1999, at URL <http://nadp.sws.uiuc.edu/nadpdata>
- Naus, C.A., 1999, Geochemistry of ground water in the Madison and Minnelusa aquifers in the Black Hills of South Dakota: Rapid City, South Dakota School of Mines and Technology, unpublished M.S. thesis, 142 p.
- Parkhurst, D.L., 1995, User's guide to PHREEQC—a computer program for speciation, reaction-path, advective-transport, and inverse geochemical calculations: U.S. Geological Survey Water-Resources Investigations Report 95-4227, 143 p.
- Parkhurst, D.L., and Appelo, C.A.J., 1999, User's guide to PHREEQC (version 2)—A computer program for speciation, batch-reaction, one-dimensional transport, and inverse geochemical calculations: U.S. Geological Survey Water-Resources Investigations Report 99-4259, 312 p.
- Peter, K.D., 1985, Availability and quality of water from the bedrock aquifers in the Rapid City area, South Dakota: U.S. Geological Survey Water-Resources Investigations Report 85-4022, 34 p.
- Plummer, L.N., Busby, J.F., Lee, R.W., and Hanshaw, B.B., 1990, Geochemical modeling of the Madison aquifer in parts of Montana, Wyoming, and South Dakota: *Water Resources Research*, v. 26, no. 9, p. 1981-2014.
- Rahn, P.H., 1971, The hydrologic significance of the November 1968 dye test on Boxelder Creek, Black Hills, South Dakota: *Proceedings of the South Dakota Academy of Science*, v. 50, p. 52-56.
- 1985, Ground water stored in the rocks of western South Dakota, *in* Rich, F.J., ed., *Geology of the Black Hills, South Dakota and Wyoming* (2d ed.): Geological Society of America, Field Trip Guidebook, American Geological Institute, p. 154-174.
- Rahn, P.H., and Gries, J.P., 1973, Large springs in the Black Hills, South Dakota and Wyoming: *South Dakota Geological Survey Report of Investigations* 107, 46 p.
- Reilly, T.E., Plummer, L.N., Phillips, P.J., and Busenberg, E., 1994, The use of simulation and multiple environmental tracers to quantify groundwater flow in a shallow aquifer: *Water Resources Research*, v. 30, no. 2, p. 421-434.
- Richter, J., Szymczak, P., Abraham, T., and Jordan, H., 1993, Use of combinations of lumped-parameter models to interpret groundwater isotope data: *Journal of Contaminant Hydrology*, v. 14, p. 1-13.
- Rose, S., 1993, Environmental tritium systematics of baseflow in Piedmont Province watersheds, Georgia: *Journal of Hydrology*, v. 143, p. 191-216.
- Stiff, H.A., Jr., 1951, The interpretation of chemical water analysis by means of patterns: *Journal of Petroleum Technology*, v. 3, no. 10, p. 15-17.
- Strobel, M.L., Galloway, J.M., Hamade, G.R., and Jarrell, G.J., 2000a, Potentiometric surface of the Madison aquifer, Black Hills area, South Dakota: U.S. Geological Survey Hydrologic Investigation Atlas HA-745-D, 2 sheets, scale 1:100,000.

- 2000b, Potentiometric surface of the Minnelusa aquifer, Black Hills area, South Dakota: U.S. Geological Survey Hydrologic Investigation Atlas HA-745-C, 2 sheets, scale 1:100,000.
- Strobel, M.L., Jarrell, G.J., Sawyer, J.F., Schleicher, J.R., and Fahrenbach, M.D., 1999, Distribution of hydrogeologic units in the Black Hills area, South Dakota: U.S. Geological Survey Hydrologic Investigations Atlas HA-743, 3 sheets, scale 1:100,000.
- Swenson, F.A., 1968, New theory of recharge to the artesian basin of the Dakotas: Geological Society of America Bulletin 1081-B, p. 163-182.
- U.S. Department of Commerce, 1999, Climatological data for South Dakota, annual summary: Asheville, N.C.
- U.S. Geological Survey, 2000, Water resources data, South Dakota, water year 1999: U.S. Geological Survey Water-Data Report SD-99-1, 501 p.
- von Morlot, A., 1847, Veber Dolomit und seine kunstliche Darstellung aus kalkstein: Naturwissenschaftliche Abhandlungenm, gesammelt und durch Subscription lursg, von Wilhelm Haidinger, v. 1, p. 305-315.
- Wenker, G.A., 1997, Geological setting and water quality of headwater springs in the Black Hills of South Dakota: Rapid City, South Dakota School of Mines and Technology, M.S. thesis, 101 p.
- Whalen, P.J., 1994, Source aquifers for Cascade Springs, Hot Springs, and Beaver Creek Springs in the southern Black Hills of South Dakota: Rapid City, South Dakota School of Mines and Technology, unpublished M.S. thesis, 299 p.
- Whitehead, R.L., 1996, Ground water atlas of the United States—segment 8, Montana, North Dakota, South Dakota, Wyoming: U.S. Geological Survey Hydrologic Investigations Atlas 730-I, 24 p.
- Yurtsever, Y., and Payne, B.R., 1986, Mathematical models based on compartmental simulation approach for quantitative interpretation of tracer data in hydrological systems: Proceedings of the 5th International Symposium on Underground Water Tracing, v. 5, p. 341-353.
- Zuber, 1986, Mathematical models for the interpretation of environmental radioisotopes in groundwater systems, *in* Fritz, P., and Fontes, J.C., eds., Handbook of Environmental Geochemistry, v. 2, The terrestrial environment: New York, Elsevier Publishing Company, p. 1-59.